arXiv:1703.08907v1 [math.OA] 27 Mar 2017

HNN EXTENSIONS OF QUASI-LATTICE ORDERED GROUPS AND THEIR OPERATOR ALGEBRAS

ASTRID AN HUEF, IAIN RAEBURN, AND ILIJA TOLICH

Abstract. The Baumslag-Solitar group is an example of an HNN extension. Spielberg showed that it has a natural positive cone, and that it is then a quasi-lattice ordered group in the sense of Nica. We give conditions for an HNN extension of a quasi-lattice ordered group (G, P) to be quasi-lattice ordered. In that case, if (G, P) is amenable as a quasi-lattice ordered group, then so is the HNN extension.

1. Introduction

Since they were introduced by Nica [9], quasi-lattice ordered groups and their C^*-algebras have generated considerable interest (see, for example, [5],[6]). The amenability of quasi-lattice ordered groups has been a deep subject (see, for example, [3],[4] and [7]). Quasi-lattice ordered groups are also examples of the more recent LCM semigroups [1], [13]. Here we generalise two recent results about the Baumslag-Solitar group.

First, Spielberg proved that the Baumslag-Solitar group is quasi-lattice ordered [11]. The Baumslag-Solitar group is an example of an HNN extension of $\mathbb{Z}$, and hence we wondered if HNN extensions could provide new classes of quasi-lattice ordered groups. Spielberg also showed that a groupoid associated to the Baumslag-Solitar semigroup is amenable [11, Theorem 3.22].

Second, Clark, an Huef and Raeburn examined the phase-transitions of the Toeplitz algebra of the Baumslag-Solitar group [2]. As part of their investigation they proved that the Baumslag-Solitar group is amenable as a quasi-lattice ordered group. The standard way to prove amenability, introduced by Laca and Raeburn [5], is to use a "controlled map": an order-preserving homomorphism between quasi-lattice ordered groups. They observed that the height map, which counts the number of times the stable letter of the HNN extension appears in a word, is almost a controlled map, and then they adapted the standard proof in [2, Appendix A] to fit.

Our innovation in this paper is a more general definition of a controlled map. We prove in Theorem 3.2 that if (G, P) is a quasi-lattice ordered group and there is a controlled map μ into an amenable group, and if $\ker \mu$ is an amenable quasi-lattice ordered group, then G is amenable. The motivation for Theorem 3.2 was two-fold. First, if a normal subgroup N of a group G is amenable and G/N is amenable, then G is amenable, and second, Spielberg's result on amenability of groupoids [12, Proposition 9.3].

Date: 8 March, 2017.

2010 *Mathematics Subject Classification.* 46L55, 46L05.

Key words and phrases. Toeplitz algebras, quasi-lattice order, HNN extension, Baumslag-Solitar groups, amenability.

This research was supported by the Marsden Fund of the Royal Society of New Zealand and by a University of Otago Publishing Bursary.

In Theorem 4.1 we give conditions under which an HNN extension of a quasi-lattice ordered group is quasi-lattice ordered. This result allows us to construct many new examples of quasi-lattice ordered groups. Finally, we use Theorem 3.2 to prove that an HNN extension of an amenable quasi-lattice ordered group is amenable (Theorem 5.1).

2. Preliminaries

Let P be a subsemigroup of a discrete group G such that $P \cap P^{-1} = \{e\}$. There is a partial order on G defined by

$$x \leq y \Leftrightarrow x^{-1}y \in P \Leftrightarrow y \in xP.$$

The order is left-invariant in the sense that $x \leq y$ implies $zx \leq zy$ for all $z \in G$. A partially ordered group (G, P) is *quasi-lattice ordered* if every finite subset of G with a common upper bound in P has a least common upper bound in P [9, Definition 2.1]. By [3, Lemma 7], (G, P) is quasi-lattice ordered if and only if:

(2.1) if $x \in PP^{-1}$, then there exist a pair $\mu, \nu \in P$ with $x = \mu\nu^{-1}$ such that $\gamma, \delta \in P$ and $\gamma\delta^{-1} = \mu\nu^{-1}$ imply $\mu \leq \gamma$ and $\nu \leq \delta$. (The pair μ, ν is unique.)

Let (G, P) be a quasi-lattice ordered group, and let $x, y \in G$. If x and y have a common upper bound in P, then their least common upper bound in P is denoted $x \vee y$. We write $x \vee y = \infty$ when x and y have no common upper bound in P and $x \vee y < \infty$ when they have a common upper bound. An *isometric representation of* P in a C^*-algebra A is a map $W : P \to A$ such that $W_e = 1$, W_p is an isometry and $W_pW_q = W_{pq}$ for all $p, q \in P$. We say that W is *covariant* if

$$W_pW_p^*W_qW_q^* = \begin{cases} W_{p\vee q}W_{p\vee q}^* & \text{if } p \vee q < \infty \\ 0 & \text{otherwise.} \end{cases} \tag{2.2}$$

Equivalently, W is covariant if

$$W_p^*W_q = \begin{cases} W_{p^{-1}(p\vee q)}W_{q^{-1}(p\vee q)}^* & \text{if } p \vee q < \infty \\ 0 & \text{otherwise.} \end{cases}$$

An example of a covariant representation is $T : P \to B(\ell^2(P))$ characterised by $T_p\epsilon_x = \epsilon_{px}$ where $\{\epsilon_x : x \in P\}$ is the orthonormal basis of point masses in $\ell^2(P)$.

In [9, §§2.4 and 4.1] Nica examined two C^*-algebras associated to (G, P). The reduced C^*-algebra $C_r^*(G, P)$ of (G, P) is the C^*-subalgebra of $B(\ell^2(P))$ generated by $\{T_p : p \in P\}$. The universal C^*-algebra $C^*(G, P)$ of (G, P) is generated by a universal covariant representation w; it is universal for covariant representations of P in the following sense: for any covariant representation $W : P \to A$ there exists a unital homomorphism $\pi_W : C^*(G, P) \to A$ such that $\pi_W(w_p) = W_p$. It follows from (2.2) that

$$C^*(G, P) = \overline{\text{span}}\{w_pw_q^* : p, q \in P\}.$$

Nica defined (G, P) to be *amenable* if the homomorphism $\pi_T : C^*(G, P) \to C_r^*(G, P)$ is faithful [9, §4.2]. He identified an equivalent condition: there exists a conditional expectation $E : C^*(G, P) \to \overline{\text{span}}\{w_pw_p^* : p \in P\}$, and (G, P) is amenable if and only if E is faithful (that is, $E(a^*a) = 0$ implies $a^*a = 0$ for all $a \in C^*(G, P)$). Laca and Raeburn took this second condition as their definition of amenability [5, Definition 3.4].

3. Order-preserving maps and amenability

A key technique, introduced by Laca and Raeburn in [5, Proposition 6.6][1], is the use of an order-preserving homomorphism between two quasi-lattice ordered groups which preserves the least upper bound structure. Crisp and Laca called such a homomorphism a *controlled map* [4]. If (G, P) and (K, Q) are quasi-lattice ordered groups, $\mu : G \to K$ is a controlled map and K is an amenable group, then (G, P) is amenable as a quasi-lattice ordered group by [5, Proposition 6.6]. Motivated by work in [2, Appendix A] we now give a weaker definition for a controlled map. We then follow the program of [2] to generalise [5, Proposition 6.6]. We state this generalisation in Theorem 3.2 below; its proof will take up the remainder of this section.

Definition 3.1. Let (G, P) and (K, Q) be quasi-lattice ordered groups. Let $\mu : G \to K$ be an order-preserving group homomorphism. For each $k \in Q$, let Σ_k be the set of $\sigma \in \mu^{-1}(k) \cap P$ which are minimal in the sense that

$$x \in \mu^{-1}(k) \cap P \text{ and } x \leq \sigma \Rightarrow \sigma = x.$$

We say μ is a *controlled map* if it has the following properties:

(1) For all $x, y \in G$ such that $x \vee y < \infty$ we have $\mu(x) \vee \mu(y) = \mu(x \vee y)$.
(2) For all $k \in Q$, Σ_k is complete in the following sense: for every $x \in \mu^{-1}(k) \cap P$ there exists $\sigma \in \Sigma_k$ such that $\sigma \leq x$.
(3) For all $k \in Q$ and $\sigma, \tau \in \Sigma_k$ we have $\sigma \vee \tau < \infty \Rightarrow \sigma = \tau$.

Theorem 3.2. *Let (G, P) and (K, Q) be quasi-lattice ordered groups. Suppose that $\mu : G \to K$ is a controlled map. If K is an amenable group and $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is an amenable quasi-lattice ordered group, then (G, P) is amenable.*

We start by showing that the kernel of a controlled map is a quasi-lattice ordered group.

Lemma 3.3. *Let (G, P) and (K, Q) be quasi-lattice ordered groups, and suppose that $\mu : G \to K$ is a controlled map. Then $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is a quasi-lattice ordered group.*

Proof. It is clear that $\mu^{-1}(e)$ is a subgroup of G and that$\mu^{-1}(e) \cap P$ is a unital semigroup. Suppose that $x, y \in \mu^{-1}(e)$ have a common upper bound $z \in \mu^{-1}(e) \cap P$. We know that z is a common upper bound for x, y in P, and hence $x \vee y$ exists in P and $x \vee y \leq z$. Now $\mu(x \vee y) = \mu(x) \vee \mu(y) = e$, and hence $x \vee y \in \mu^{-1}(e) \cap P$. Thus $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is a quasi-lattice ordered group. □

To prove Theorem 3.2 we will show that the conditional expectation

$$E : C^*(G, P) \to \overline{\operatorname{span}}\{w_p w_p^* : p \in P\}$$

is faithful. We will use the amenability of K to construct a faithful conditional expectation $\Psi^\mu : C^*(G, P) \to \overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$, and then show that E is faithful when restricted to range Ψ^μ. To construct Ψ^μ we follow the method of [5, Lemma 6.5] which uses a coaction.

[1] There is an error in the statement of [5, Proposition 6.6]: the final line should read "if $\mathcal{G}$ is amenable then (G, P) is amenable".

Let G be a discrete group and let A be a unital C^*-algebra. Let

$$\delta_G : C^*(G) \to C^*(G) \otimes_{\min} C^*(G)$$

be the comultiplication of G which is characterised by $\delta_G(u_g) = u_g \otimes u_g$ for $g \in G$. A *coaction* of G on A is a unital homomorphism $\delta : A \to A \otimes_{\min} C^*(G)$ such that

$$(\delta \otimes \mathrm{id}) \circ \delta = (\mathrm{id} \otimes \delta_G) \circ \delta.$$

We say that δ is *nondegenerate* if $\delta(A)(1 \otimes C^*(G)) = A \otimes_{\min} C^*(G)$.

Lemma 3.4. *Let (G, P) be a quasi-lattice ordered group. Suppose that there exists a group K and a homomorphism $\mu : G \to K$. Then there exists an injective coaction*

$$\delta_\mu : C^*(G, P) \to C^*(G, P) \otimes_{\min} C^*(K)$$

characterised by $\delta_\mu(w_p) = w_p \otimes u_{\mu(p)}$ for all $p \in P$.

Proof. Let $W : P \to C^*(G, P) \otimes_{\min} C^*(K)$ be characterised by $W_p = w_p \otimes u_{\mu(p)}$. We will show that W is a covariant representation, and then take $\delta_\mu := \pi_W$. Unitaries are isometries and hence W_p is isometric for all $p \in P$. Observe that $W_e = w_e \otimes u_{\mu(e)} = 1 \otimes 1$, and

$$W_p W_q = w_p w_q \otimes u_{\mu(p)} u_{\mu(q)} = w_{pq} \otimes u_{\mu(pq)} = W_{pq} \text{ for all } p, q \in P.$$

Thus W is an isometric representation. To prove W is covariant, we fix $x, y \in P$ and compute:

$$\begin{aligned} W_x W_x^* W_y W_y^* &= w_x w_x^* w_y w_y^* \otimes u_{\mu(x)} u_{\mu(x)}^* u_{\mu(y)} u_{\mu(y)}^* \\ &= \begin{cases} w_{x\vee y} w_{x\vee y}^* \otimes 1 & \text{if } x \vee y < \infty \\ 0 \otimes 1 & \text{otherwise} \end{cases} \\ &= \begin{cases} w_{x\vee y} w_{x\vee y}^* \otimes w_{\mu(x\vee y)} w_{\mu(x\vee y)}^* & \text{if } x \vee y < \infty \\ 0 & \text{otherwise} \end{cases} \\ &= W_{x\vee y} W_{x\vee y}^*. \end{aligned}$$

Thus W is a covariant representation of P. By the universal property of $C^*(G, P)$, there exists a homomorphism $\delta_\mu := \pi_W$, which has the desired properties. Since $W_e = 1 \otimes 1$ it follows that δ_μ is unital.

To prove the comultiplication identity, we compute on generators: for $p, q \in P$ we have

$$\begin{aligned} ((\delta_\mu \otimes \mathrm{id}) \circ \delta_\mu)(w_p w_q^*) &= (\delta_\mu \otimes \mathrm{id})(w_p w_q^* \otimes u_{\mu(pq^{-1})}) \\ &= \delta_\mu(w_p w_q^*) \otimes \mathrm{id}(u_{\mu(pq^{-1})}) \\ &= w_p w_q^* \otimes u_{\mu(pq^{-1})} \otimes u_{\mu(pq^{-1})} \\ &= w_p w_q^* \otimes \delta_K(u_{\mu(pq^{-1})}) \\ &= \mathrm{id} \otimes \delta_K(w_p w_q^* \otimes u_{\mu(pq^{-1})}) \\ &= ((\mathrm{id} \otimes \delta_K) \circ \delta_\mu)(w_p w_q^*). \end{aligned}$$

Hence $(\delta_\mu \otimes \mathrm{id}) \circ \delta_\mu = (\mathrm{id} \otimes \delta_K) \circ \delta_\mu$. Thus δ_μ is a coaction.

To show that δ_μ is injective, let $\pi : C^*(G, P) \to B(H)$ be a faithful representation. We will show that π can be written as a composition of δ_μ and another representation.

Let $\epsilon : C^*(K) \to \mathbb{C}$ be the trivial representation on $\mathbb{C}$ such that $\epsilon(u_k) = 1$ for all $k \in K$. By the properties of the minimal tensor product (see [10, Proposition B.13]) there exists a homomorphism

$$\pi \otimes \epsilon : C^*(G,P) \otimes_{\min} C^*(G) \to B(H) \otimes \mathbb{C} = B(H).$$

Since

$$(\pi \otimes \epsilon) \circ \delta_\mu(w_p) = (\pi \otimes \epsilon)(w_p \otimes u_{\mu(p)}) = \pi(w_p),$$

we have $\pi = (\pi \otimes \epsilon) \circ \delta_\mu$. Now suppose that $\delta_\mu(a) = 0$ for some $a \in C^*(G,P)$. Then $0 = (\pi \otimes \epsilon) \circ \delta_\mu(a) = \pi(a)$. Since π is faithful, $a = 0$. Hence δ_μ is injective.

To prove that δ_μ is a nondegenerate coaction we must show that

$$\delta_\mu(C^*(G,P))(1 \otimes C^*(K)) = C^*(G,P) \otimes_{\min} C^*(K).$$

It suffices to show that we can get the spanning elements $w_p w_q^* \otimes u_k$, and this is easy:

$$\delta_\mu(w_p w_q^*)(1 \otimes u_{\mu(qp^{-1})k}) = w_p w_q^* \otimes u_{\mu(pq^{-1})}(1 \otimes u_{\mu(qp^{-1})k}) = w_p w_q^* \otimes u_k.$$

Thus δ_μ is nondegenerate. □

Let λ be the left-regular representation of a discrete group K. There is a trace τ on $C^*(K)$ characterised by

$$\tau(u_k) = (\lambda_k \epsilon_e \mid \epsilon_e) = \begin{cases} 1 & \text{if } k = e \\ 0 & \text{otherwise.} \end{cases}$$

It is well-known that if K is an amenable group, then τ is faithful.

Lemma 3.5. *Let (G,P) be a quasi-lattice ordered group. Suppose that there exist a group K and a homomorphism $\mu : G \to K$. Let*

$$\delta_\mu : C^*(G,P) \to C^*(G,P) \otimes_{\min} C^*(K)$$

be the coaction of Lemma 3.4. Then

$$\Psi^\mu := (\mathrm{id} \otimes \tau) \circ \delta_\mu$$

is a conditional expectation of $C^*(G,P)$ with range $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$. If K is an amenable group, then Ψ^μ is faithful.

Proof. Since $\mathrm{id} \otimes \tau$ and δ_μ are linear and norm decreasing, so is Ψ^μ. Since $\Psi^\mu(w_e) = 1$ the norm of Ψ^μ is 1. We have

$$\Psi^\mu(w_p w_q^*) = \begin{cases} w_p w_q^* & \text{if } \mu(p) = \mu(q) \\ 0 & \text{otherwise,} \end{cases} \tag{3.1}$$

and hence $\Psi^\mu \circ \Psi^\mu = \Psi^\mu$. Thus Ψ^μ is a conditional expectation by [14].

From (3.1) we see that $\overline{\operatorname{span}}\{w_p w_q^* : \mu(q) = \mu(p)\} \subseteq \operatorname{range} \Psi^\mu$. To show the reverse inclusion, fix $b \in \operatorname{range} \Psi^\mu$, say $b = \Psi^\mu(a)$ for some $a \in C^*(G,P)$. Also fix $\epsilon > 0$. There exists a finite subset $F \subseteq P \times P$ such that $\|a - \sum_{(p,q)\in F} \lambda_{p,q} w_p w_q^*\| < \epsilon$. Since Ψ^μ is linear and norm-decreasing,

$$\epsilon > \Big\| a - \sum_{(p,q)\in F} \lambda_{p,q} w_p w_q^* \Big\| \geq \Big\| \Psi^\mu\Big(a - \sum_{(p,q)\in F} \lambda_{p,q} w_p w_q^* \Big) \Big\|$$

$$= \Big\| \Psi^\mu(a) - \Psi^\mu\Big(\sum_{(p,q)\in F} \lambda_{p,q} w_p w_q^* \Big) \Big\| = \Big\| b - \sum_{(p,q)\in F,\ \mu(p)=\mu(q)} \lambda_{p,q} w_p w_q^* \Big\|.$$

Thus $b \in \overline{\operatorname{span}}\{w_p w_q^* : \mu(q) = \mu(p)\}$, and $\operatorname{range}\Psi^\mu = \overline{\operatorname{span}}\{w_p w_q^* : \mu(q) = \mu(p)\}$.

Now suppose that K is amenable. To see that Ψ^μ is faithful, we follow the proof of [5, Lemma 6.5]. Let $a \in C^*(G,P)$ and suppose that $\Psi^\mu(a^*a) = 0$. Let f be an arbitrary state on $C^*(G,P)$. Then

$$\begin{aligned} 0 &= f(\Psi^\mu(a^*a)) = f \circ (\mathrm{id} \otimes \tau) \circ \delta_\mu(a^*a) \\ &= (f \otimes \tau) \circ \delta_\mu(a^*a) = \tau \circ (f \otimes \mathrm{id}) \circ \delta_\mu(a^*a). \end{aligned}$$

Since K is amenable, τ is faithful. Hence $(f \otimes \mathrm{id}) \circ \delta(a^*a) = 0$. This implies that for all states f on $C^*(G,P)$ and states g on $C^*(K)$,

$$g \circ (f \otimes \mathrm{id}) \circ \delta_\mu(a^*a) = (f \otimes g) \circ \delta_\mu(a^*a) = 0.$$

To see that $\delta_\mu(a^*a) = 0$, let $\pi_1 : C^*(G,P) \to H_1$ and $\pi_2 : C^*(K) \to H_2$ be faithful representations. Then $\pi_1 \otimes \pi_2$ is a faithful representation of $C^*(G,P) \otimes_{\min} C^*(K)$ on $B(H_1 \otimes H_2)$ by [10, Corollary B.11]. Fix unit vectors $h \in H_1$, $k \in H_2$. There exists a state $f_h \otimes f_k$ on $C^*(G,P) \otimes_{\min} C^*(K)$ defined by

$$f_h \otimes f_k(x) = (\pi_1 \otimes \pi_2(x)(h \otimes k) \mid h \otimes k).$$

Since $(f \otimes g) \circ \delta_\mu(a^*a) = 0$ for all states f of $C^*(G,P)$ and g of $C^*(K)$, we have

$$\begin{aligned} 0 &= f_h \otimes f_k(\delta_\mu(a^*a)) \\ &= (\pi_1 \otimes \pi_2(\delta_\mu(a^*a))(h \otimes k) \mid h \otimes k) \\ &= (\pi_1 \otimes \pi_2(\delta_\mu(a))(h \otimes k) \mid \pi_1 \otimes \pi_2(\delta_\mu(a))h \otimes k) \\ &= \|\pi_1 \otimes \pi_2(\delta_\mu(a))(h \otimes k)\|^2. \end{aligned}$$

Hence $\pi_1 \otimes \pi_2(\delta_\mu(a^*a)) = 0$. Since $\pi_1 \otimes \pi_2$ is faithful, $\delta_\mu(a^*a)) = 0$. But δ_μ is injective, and hence $a = 0$, and Ψ^μ is faithful. □

Next we investigate the structure of

$$\operatorname{range}\Psi^\mu = \overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}.$$

In the statement of the next lemma, note that we can view $C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ as a C^*-subalgebra of $C^*(G,P)$.

Lemma 3.6. *Let (G,P) and (K,Q) be quasi-lattice ordered groups, and suppose that $\mu : G \to K$ a controlled map. Let $k \in Q$, and let F be a finite subset of Σ_k. Let*

$$B_{k,F} := \operatorname{span}\{w_\sigma D w_\tau^* : \sigma, \tau \in F, D \in C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)\}.$$

Then $B_{k,F}$ is a closed C^*-subalgebra of $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$.

Proof. To see that $B_{k,F}$ is contained in $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$, it suffices to consider D of the form $w_\alpha w_\beta^*$ where $\mu(\alpha) = \mu(\beta) = e$. Then $w_\sigma D w_\tau^* = w_{\sigma\alpha} w_{\tau\beta}^*$ and $\mu(\sigma\alpha) = \mu(\sigma) = k = \mu(\tau) = \mu(\tau\beta)$. Thus $B_{k,F}$ is a subalgebra of $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$.

We will prove the lemma by showing that $B_{k,F}$ is isomorphic to

$$M_F(\mathbb{C}) \otimes C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big).$$

By Definition 3.1(3), the elements of F have no common upper bound unless they are equal. So

$$(w_\sigma w_\tau^*)(w_\gamma w_\delta^*) = w_{\sigma\tau^{-1}(\tau\vee\gamma)} w^*_{\delta\gamma^{-1}(\tau\vee\gamma)} = \begin{cases} w_\sigma w_\delta^* & \text{if } \tau = \gamma \\ 0 & \text{otherwise.} \end{cases}$$

Thus $\{w_\sigma w_\tau^* : \sigma, \tau \in F\}$ is a set of matrix units in the C^*-algebra $\overline{B_{k,F}}$. This gives a homomorphism $\theta : M_F(\mathbb{C}) \to \overline{B_{k,F}}$ which maps the matrix units $\{E_{\sigma,\tau} : \sigma, \tau \in F\}$ in $M_F(\mathbb{C})$ to $\{w_\sigma w_\tau^* : \sigma, \tau \in F\} \subset B_{k,F}$. It is easy to check that the formula

$$\psi(D) = \sum_{\gamma \in F} w_\gamma D w_\gamma^*$$

gives a homomorphism $\psi : C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big) \to B_{k,F}$. We have

$$\begin{aligned} \theta(E_{\sigma,\tau})\psi(D) &= w_\sigma w_\tau^* \sum_{\gamma \in F} w_\gamma D w_\gamma^* \\ &= w_\sigma w_\tau^* w_\tau D w_\tau^* \quad (w_\tau^* w_\gamma = 0 \text{ unless } \tau = \gamma) \\ &= w_\sigma D w_\tau^* \\ &= (\sum_{\gamma \in F} w_\gamma D w_\gamma^*) w_\sigma w_\tau^* \\ &= \psi(D)\theta(E_{\sigma,\tau}). \end{aligned}$$

Each $A \in M_F(\mathbb{C})$ is a linear combination of the $E_{\sigma,\tau}$, and hence $\psi(D)\theta(A) = \theta(A)\psi(D)$ for all $A \in M_F(\mathbb{C})$ and $D \in C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$. Since the ranges of θ and ψ commute, the universal property of the maximal tensor product gives a homomorphism $\theta \otimes_{\max} \psi$ of $M_F(\mathbb{C}) \otimes_{\max} C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ into $\overline{B_{k,F}}$.

By [10, Theorem B.18]

$$\begin{aligned} &M_F(\mathbb{C}) \otimes_{\max} C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big) \\ &\qquad = \operatorname{span}\{E_{\sigma,\tau} \otimes D : \sigma, \tau \in F \text{ and } D \in C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)\}, \end{aligned}$$

with no closure. So the range of $\theta \otimes_{\max} \psi$ is spanned by $\theta(E_{\sigma,\tau})\psi(D) = w_\sigma D w_\tau^*$ and hence is $B_{k,F}$. Thus $B_{k,F}$ is a closed C^*-subalgebra of $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$. □

Let $\{\epsilon_x : x \in P\}$ be the usual basis for $\ell^2(P)$. Let T be the covariant representation of (G, P) on $\ell^2(P)$ such that $T_p \epsilon_x = \epsilon_{px}$, and let π_T be the corresponding homomorphism of $C^*(G, P)$ onto $C_r^*(G, P)$ such that $\pi_T(w_p) = T_p$. For $k \in Q$ we consider the subspaces

$$H_k := \{\epsilon_{\gamma z} : \gamma \in \Sigma_k, z \in \mu^{-1}(e) \cap P\}.$$

Lemma 3.7. *Let (G, P) and (K, Q) be quasi-lattice ordered groups, and suppose that $\mu : G \to K$ is a controlled map. Let $k \in Q$. Then*

1. *H_k is invariant for $\pi_T|_{B_{k,F}}$;*
2. *if $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable, then $\pi_T|_{B_{k,F}}$ is isometric on H_k.*

Proof. For (1), let $\sigma, \tau \in F$ and let $x, y \in \mu^{-1}(e)$ and let $\epsilon_{\gamma z} \in H_k$. Then $w_\sigma w_x w_y^* w_\tau^*$ is a spanning element of $B_{k,F}$. Since $\mu(\tau) = k = \mu(\gamma)$ we have

$$\pi_T(w_\sigma w_x w_y^* w_\tau^*)\epsilon_{\gamma z} = \begin{cases} \epsilon_{\sigma x y^{-1} z} & \text{if } \gamma = \tau \text{ and } y \le z \\ 0 & \text{otherwise.} \end{cases}$$

If $\pi_T(w_\sigma w_x w_y^* w_\tau^*)\epsilon_{\gamma z} = 0$ we are done. Otherwise, to see that $\epsilon_{\sigma x y^{-1} z}$ is back in H_k, suppose that $y \le z$. Then $y^{-1}z \in P$. Since $\sigma x \in \mu^{-1}(k) \cap P$ we have $\epsilon_{(\sigma x)(y^{-1}z)} \in H_k$. It follows that H_k is invariant for $\pi_T|_{B_{k,F}}$.

For (2) suppose that $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable. We will show that $\pi_T|_{B_{k,F}}$ is faithful on H_k. Take $B = \sum_{\sigma,\tau \in F} w_\sigma D_{\sigma,\tau} w_\tau^* \in B_{k,F}$ and suppose that $\pi_T(B)|_{H_k} = 0$. Fix $\gamma, \delta \in F$. Then

$$T_\gamma^* \pi_T(B) T_\delta = \pi_T(w_\gamma^*)\pi_T(B)\pi_T(w_\delta) = \pi_T(D_{\gamma,\delta})$$

Since T_δ is an injection from H_e to H_k and $\pi_T(B)|_{H_k} = 0$, it follows that $\pi_T(B)T_\delta|_{H_e} = 0$. Thus $\pi_T(D_{\gamma,\delta})|_{H_e} = 0$.

But the restriction

$$(\pi_T|_{C^*\left(\mu^{-1}(e), \mu^{-1}(e)\cap P\right)})|_{H_e}$$

is the Toeplitz representation of $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$, and hence is faithful by amenability. Thus $D_{\gamma,\delta} = 0$. Repeating the argument finitely many times shows that all the $D_{\sigma,\tau} = 0$ and hence that $B = 0$. Thus $\pi_T|_{B_{k,F}}$ is faithful on H_k, and therefore is isometric. □

Lemma 3.8. *Let (G, P) and (K, Q) be quasi-lattice ordered groups, and suppose that $\mu : G \to K$ is a controlled map. Let $B_k = \overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q) = k\}$. Let $\mathcal{F}$ be the set of all finite sets $F \subseteq \Sigma_k$. Then $B_k = \overline{\cup_{F\in\mathcal{F}} B_{k,F}}$. If $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable, then $\pi_T|_{B_k}$ is isometric on H_k.*

Proof. Observe that $\mathcal{F}$ is a directed set partially ordered by inclusion with $E, F \in \mathcal{F}$ majorismajorizeded by $E \cup F$. If $E \subseteq F$, then $B_{k,E} \subseteq B_{k,F}$. Thus $\{B_{k,F} : F \in \mathcal{F}\}$ is an inductive system with limit $\overline{\cup_{F\in\mathcal{F}} B_{k,F}}$.

For each $F \in \mathcal{F}$ we have $B_{k,F} \subseteq B_k$, and B_k is closed. Therefore $\overline{\cup_{F\in\mathcal{F}} B_{k,F}} \subseteq B_k$. To prove the reverse inclusion it suffices to show that the spanning elements of B_k are in $B_{k,F}$ for some F. Fix $p, q \in P$ such that $\mu(p) = \mu(q) = k$ and consider $w_p w_q^*$. By Definition 3.1(2), the set Σ_k of minimal elements is complete, and there exists $\sigma, \tau \in \Sigma_k$ such that $\sigma \le p$ and $\tau \le q$. Hence there exists $x, y \in P$ such that $p = \sigma x$ and $q = \tau y$. Thus $w_p w_q^* = w_{\sigma x} w_{\tau y}^* = w_\sigma (w_x w_y^*) w_\tau^*$ and $w_x w_y^* \in C^*\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$. Since $\{\sigma, \tau\} \in \mathcal{F}$ we have $w_p w_q^* \in B_{k,\{\sigma,\tau\}}$. Thus $B_k \subseteq \overline{\cup_{F\in\mathcal{F}} B_{k,F}}$, and equality follows.

Finally, suppose that $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable. Then $\pi_T|_{B_{k,F}}$ is isometric on H_k for all $F \in \mathcal{F}$ by Lemma 3.7(2). Since π_T is isometric on every $B_{k,F}$, its extension to the closure is also isometric. □

Let $\mathcal{I}$ be the set of all finite sets $I \subset Q$ that are closed under $\vee$ in the sense that $s, t \in I$ and $s \vee t < \infty$ implies that $s \vee t \in I$.

Lemma 3.9. *Let (G,P) and (K,Q) be quasi-lattice ordered groups, and suppose that $\mu : G \to K$ is a controlled map. For each $I \in \mathcal{I}$ let*

$$C_I = \overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q) \in I\}.$$

Then C_I is a C^*-subalgebra of $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$, $C_I = \operatorname{span}\{B_k : k \in I\}$ and $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\} = \overline{\cup_{I\in\mathcal{I}} C_I}$.

Proof. Fix $I \in \mathcal{I}$. To see that C_I is a C^*-subalgebra, it suffices to show that $\operatorname{span}\{w_p w_q^* : \mu(p) = \mu(q) \in I\}$ is a $*$-subalgebra. It's clearly closed under taking adjoints. Let $p,q,r,s \in P$ such that $\mu(p) = \mu(q) \in I$ and $\mu(r) = \mu(s) \in I$. Then

$$w_p w_q^* w_r w_s^* = \begin{cases} w_{pq^{-1}(q\vee r)} w^*_{sr^{-1}(q\vee r)} & \text{if } q \vee r < \infty \\ 0 & \text{otherwise.} \end{cases}$$

If $w_p w_q^* w_r w_s^* = 0$ we are done. So suppose that $w_p w_q^* w_r w_s^* \neq 0$. Then $q \vee r < \infty$. Since μ is a controlled map and $\mu(p) = \mu(q)$, by Definition 3.1(1),

$$\mu(pq^{-1}(q \vee r)) = \mu(q \vee r) = \mu(q) \vee \mu(r).$$

Similarly, $\mu(sr^{-1}(q\vee r)) = \mu(q)\vee\mu(r)$. Since I is closed under $\vee$ we have $\mu(q)\vee\mu(r) \in I$, and hence $w_p w_q^* w_r w_s^* \in \operatorname{span}\{w_p w_q^* : \mu(p) = \mu(q) \in I\}$. It follows that C_I is a C^*-subalgebra.

For each $k \in I$, we have $B_k \subseteq C_I$, and so $\operatorname{span}\{B_k : k \in I\} \subseteq C_I$. To show the reverse inclusion observe that for $w_p w_q^* \in C_I$ we have $w_p w_q^* \in B_{\mu(p)}$. Since the finite span of closed subalgebras is closed, $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q) \in I\} \subseteq \operatorname{span}\{B_k : k \in I\}$. Thus $C_I = \operatorname{span}\{B_k : k \in I\}$. □

Proposition 3.10. *Let (G,P) and (K,Q) be quasi-lattice ordered groups, and suppose that $\mu : G \to K$ is a controlled map. If $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable, then π_T is faithful on $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\}$.*

Proof. By Lemma 3.9, $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\} = \overline{\cup_{I\in\mathcal{I}} C_I}$. Thus it suffices to show that π_T is isometric on each C_I. Fix $I \in \mathcal{I}$. Suppose that $\pi_T(R) = 0$ for some $R \in C_I$. Then there exist $R_k \in B_k$ such that $R = \sum_{k\in I} R_k$ and then $\sum_{k\in I} \pi_T(R_k) = 0$.

We claim that if $k \not\leq j$, then $\pi_T(B_k)|_{H_j} = 0$ (it then follows that $\pi_T(R_k)|_{H_j} = 0$). To prove the claim, it suffices to show that $\pi_T(w_q^*)\epsilon_{\gamma z} = 0$ for all $q \in \mu^{-1}(k) \cap P$ and $\epsilon_{\gamma z} \in H_k$. We have

$$\pi_T(w_q^*)\epsilon_{\gamma z} = T_q^*\epsilon_{\gamma z} = \begin{cases} \epsilon_{q^{-1}\gamma z} & \text{if } q \leq \gamma z \\ 0 & \text{otherwise.} \end{cases}$$

But $q \leq \gamma z$ implies $k = \mu(q) \leq \mu(\gamma z) = \mu(\gamma) = j$. So $k \not\leq j$ implies $\pi_T(w_q^*)\epsilon_{\gamma z} = 0$. Hence $\pi_T(B_k)|_{H_j} = 0$ if $k \not\leq j$ as claimed.

Let l be a minimal element of I in the sense that $x \leq l$ implies $x = l$. Then for $k \in I$, we have $k \not\leq l$ unless $k = l$. Now

$$0 = \sum_{k\in I} \pi_T(R_k)|_{H_l} = \pi_T(R_l)|_{H_l}.$$

Since $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable, $\pi_T|_{B_l}$ is isometric on H_l by Lemma 3.8. Thus $R_l = 0$.

Let l_2 be a minimal element of $I\backslash\{l\}$. Then we can repeat the above argument to get $R_{l_2} = 0$. Since I is finite, we can continue to conclude that $R = 0$. □

We can now prove Theorem 3.2

Proof of Theorem 3.2. Suppose that K is an amenable group. To see (G, P) is amenable, we will show that the conditional expectation

$$E : C^*(G, P) \to \overline{\operatorname{span}}\{w_p w_p^* : p \in P\}$$

is faithful. Let Ψ^μ be the conditional expectation of Lemma 3.5. We have

$$\begin{aligned} E(\Psi^\mu(w_p w_q^*)) &= \begin{cases} \Psi(w_p w_q^*) & \text{if } \mu(p) = \mu(q) \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} w_p w_p^* & \text{if } p = q \\ 0 & \text{otherwise} \end{cases} \\ &= \Psi(w_p w_q^*), \end{aligned}$$

and hence $E = E \circ \Psi^\mu$.

Since K is an amenable group, Ψ^μ is faithful by Lemma 3.5. Let $P_z \in B(\ell^2(P))$ be the orthogonal projection onto $\operatorname{span}\{\epsilon_z\}$. It is straightforward to show that the diagonal map $\Delta : B(\ell^2(P)) \to B(\ell^2(P))$ given by

$$\Delta(T) = \sum_{z \in P} P_z T P_z$$

is a conditional expectation such that $\Delta \circ \pi_T = \pi_T \circ E$ and is faithful.

Now suppose that $R \in C^*(G, P)$ and $E(R^*R) = 0$. Then $E(\Psi^\mu(R^*R)) = 0$ and so $\pi_T \circ E(\Psi^\mu(R^*R)) = 0$. This gives $\Delta \circ \pi_T(\Psi^\mu(R^*R)) = 0$. Since Δ is faithful, it follows that $\pi_T(\Psi^\mu(R^*R)) = 0$. Since $\big(\mu^{-1}(e), \mu^{-1}(e) \cap P\big)$ is amenable, Lemma 3.10 implies that π_T is faithful on $\overline{\operatorname{span}}\{w_p w_q^* : \mu(p) = \mu(q)\} = \operatorname{range} \Psi^\mu$. Thus $\Psi^\mu(R^*R) = 0$, and then $R = 0$ since Ψ^μ is faithful. Hence E is faithful and (G, P) is amenable. □

4. Quasi-lattice ordered HNN extensions

Let G be a group, let A and B be subgroups of G, and let $\phi : A \to B$ be an isomorphism. The group with presentation

$$G^* = \langle G, t \mid t^{-1} a t = \phi(a), a \in A \rangle$$

is the *HNN extension of* G with respect to A, B and ϕ. For every HNN extension G^* the *height map* is the homomorphism $\theta : G^* \to \mathbb{Z}$ such that $\theta(g) = 0$ for all $g \in G$ and $\theta(t) = 1$.

Example. Let $c, d \in \mathbb{Z}$. The Baumslag-Solitar group

$$\mathrm{BS}(c, d) = \langle x, t \mid t^{-1} x^d t = x^c \rangle = \langle x, t \mid t x^c = x^d t \rangle$$

is an HNN extension of $\mathbb{Z}$ with respect to $A = d\mathbb{Z}$, $B = c\mathbb{Z}$ and $\phi : A \to B$ given by $\phi(dn) = cn$ for all $n \in \mathbb{Z}$. Then $\mathbb{Z}^*$ satisfies the relation $t^{-1} d t = \phi(d) = c$. Let $\mathrm{BS}(c, d)^+$ be the subsemigroup of $\mathrm{BS}(c, d)$ generated by x and t. Spielberg showed in [11, Theorem 2.11] that $(\mathrm{BS}(c, d), \mathrm{BS}(c, d)^+)$ is quasi-lattice ordered for all $c, d > 0$; he

also proved in [11, Lemma 2.12] that $(\mathrm{BS}(c,-d), \mathrm{BS}(c,-d)^+)$ is not quasi-lattice ordered unless $c = 1$.

To work with an HNN extension we use a normal form for its elements from [8, Theorem 2.1]. We choose X to be a complete set of left coset representatives for G/A such that $xA \neq x'A$ for $x \neq x' \in X$. Similarly, choose a complete set Y of left coset representatives for G/B. Then a (right) *normal form relative to X and Y* of $g \in G$ is a product

$$g = g_0 t^{\epsilon_1} g_1 t^{\epsilon_2} \dots g_{n-1} t^{\epsilon_n} g_n$$

where:

(1) g_n is an arbitrary element of G.
(2) If $\epsilon_i = 1$, then g_{i-1} is an element of X
(3) If $\epsilon_i = -1$, then g_{i-1} is an element of Y.

By [8, Theorem 2.1], for every choice of complete left coset representatives X and Y, each $g \in G^*$ has a unique normal form.

Our goal is to generalise the properties of the Baumslag-Solitar group with $c, d > 0$ to construct quasi-lattice ordered HNN extensions of other quasi-lattice ordered groups.

Let (G, P) be a quasi-lattice ordered group. Let G^* be the HNN extension of G with respect to subgroups A and B with an isomorphism $\phi : A \to B$. Let P^* be the subsemigroup of G^* generated by P and t. In general, (G^*, P^*) is not a quasi-lattice ordered group. For example, if $c > 1$, then $(\mathrm{BS}(c,-d), \mathrm{BS}(c,-d)^+)$ is not quasi-lattice ordered by [11, Lemma 2.12]. We need some conditions on our subgroups A and B and on the isomorphism ϕ which ensure that (G^*, P^*) is quasi-lattice ordered.

There are two reasons why $(\mathrm{BS}(c,d), \mathrm{BS}(c,d)^+)$ is easy to work with. The first is that there are natural choices of coset representatives: $\{0, \dots, d-1\}$ for $A = d\mathbb{Z}$ and $\{0, \dots, c-1\}$ for $B = c\mathbb{Z}$. The second is that the subgroup isomorphism $\phi : md \mapsto mc$ takes positive elements to positive elements. In particular, using this choice of representatives, every element $\omega \in \mathrm{BS}(c,d)^+$ has a unique normal form

$$\omega = x^{m_0} t x^{m_1} t \dots x^{m_{n-1}} t x^{m_n}$$

where $0 \leq m_i < d$ for $i < n$ and $m_n \in \mathbb{N}$. This choice of coset representatives is associated to the division algorithm on $\mathbb{N}$: for every $n \in \mathbb{N}$ we can uniquely write $n = md + r$ for some $m \in \mathbb{N}$ and $0 \leq r \leq d-1$.

In general, for G^* we would like a natural choice of coset representatives for G/A and G/B so that every element of P^* has a unique normal form that is a sequence of elements in P and t.

Theorem 4.1. *Let (G, P) be a quasi-lattice ordered group with subgroups A and B. Suppose that:*

(1) *There is an isomorphism $\phi : A \to B$ such that $\phi(A \cap P) = B \cap P$;*
(2) *Every left coset $gA \in G/A$ such that $gA \cap P \neq \emptyset$ has a minimal coset representative $p \in P$: $q \in gA \cap P \Rightarrow p \leq q$;*
(3) *For every $x, y \in B$, $x \vee y < \infty \Rightarrow x \vee y \in B$.*

Let $G^* = \langle G, t \mid t^{-1}at = \phi(a), a \in A \rangle$ be the HNN extension of G and let P^* be the subsemigroup of G^* generated by $\{P, t\}$. Then (G^*, P^*) is quasi-lattice ordered.

Before we can prove Theorem 4.1, we need to prove two lemmas. The first shows that elements of P^* are guaranteed to have normal forms made up of elements of P and t if and only if condition (2) of Theorem 4.1 holds. The second is a technical lemma which we will use several times in Theorem 4.1 and in later proofs.

Lemma 4.2. *Suppose that (G, P) is a quasi-lattice ordered group with subgroups A and B. Suppose that $\phi : A \to B$ is a group isomorphism such that $\phi(A \cap P) = B \cap P$. Let $G^* = \langle G, t \mid t^{-1}at = \phi(a), a \in A\rangle$ be the corresponding HNN extension of G and let P^* be the subsemigroup of G^* generated by $P \cup \{t\}$. Let*

$$L_A := \{p \in P : q \in pA \cap P \Rightarrow p \leq q\}.$$

The following two statements are equivalent:

1. *Every left coset $gA \in G/A$ such that $gA \cap P \neq \emptyset$ has a minimal coset representative $p \in P$;*
2. *There exists a complete set X of left coset representatives such that $L_A \subseteq X$ and every $\alpha \in P^*$ has normal form*

$$\alpha = p_0 t p_1 t \dots p_{n-1} t p_n \text{ where } p_i \in L_A \text{ for all } 0 \leq i < n,\ p_n \in P. \tag{4.1}$$

Proof. Assume (1). Choose a complete set X of coset representatives which contains L_A. Let $\alpha \in P^*$. If $\theta(\alpha) = 0$ then $\alpha \in P$, and α has form (4.1) trivially.

We proceed by induction on $\theta(\alpha) \geq 1$. Suppose that $\theta(\alpha) = 1$. We may write $\alpha = q_0 t q_1$ for some $q_0, q_1 \in P$. Then $q_0 A \cap P \neq \emptyset$, and there exists $p_0 \in L_A$ such that $p_0 A = q_0 A$ and $p_0 \leq q_0$. Thus$p_0^{-1} q_0 \in P \cap A$. Hence $q_0 = p_0 a$ for some $a \in A \cap P$. Thus α has normal form

$$\alpha = p_0 a t q_1 = p_0 t \phi(a) q_1.$$

Since $\phi(A \cap P) = B \cap P$ we have $\phi(a) \in P$ and so X satisfies (2).

Suppose that all α with $1 \leq \theta(\alpha) \leq k$ have normal form (4.1). Consider α with $\theta(\alpha) = k + 1$. We write

$$\alpha = q_0 t q_1 t \dots t q_k t q_{k+1}.$$

By assumption, we can write the first $2k + 2$ terms of α in normal form

$$p_0 t p_1 t \dots p_{k-1} t r_k$$

where $p_i \in L_A$ for $0 \leq i < k$ and $r_k \in P$. There exists $p_k \in L_A$ such that $r_k A = p_k A$ and $p_k \leq r_k$. As above, we can write $r_k = p_k a$ for some $a \in A \cap P$. Then

$$\begin{aligned} \alpha &= p_0 t p_1 t \dots t r_k t q_{k+1} \\ &= p_0 t p_1 t \dots t p_k a t q_{k+1} \\ &= p_0 t p_1 t \dots t p_k t \phi(a) q_{k+1}. \end{aligned}$$

We set $p_{k+1} = \phi(a) q_{k+1}$, which is in P because $\phi(a)$ is. Then $\alpha = p_0 t p_1 t \dots t p_k t p_{k+1}$ has form (4.1). By induction, every α has normal form (4.1). This implies (2).

For (2) $\Rightarrow$ (1) we argue by contradiction: we will assume (2) holds but (1) doesn't. Let X be a set of coset representatives satisfying (2), and suppose that there exists some coset gA such that $gA \cap P \neq \emptyset$ which has no minimal coset representative. Let $p \in X$ be the coset representative of gA. Since p is not minimal, there exists $q \in gA \cap P$ with $p \not\leq q$. Thus $p^{-1} q \notin P$. Consider $qt \in P^*$ in normal form:

$$qt = p p^{-1} q t = p t \phi(p^{-1} q). \tag{4.2}$$

Since $\phi(A \cap P) = B \cap P$ we have $\phi(p^{-1}q) \notin P$. So (4.2) is not the normal form (4.1), a contradiction. Thus (2) $\Rightarrow$ (1). □

Lemma 4.3. *Let (G, P) be a quasi-lattice ordered group and let B be a subgroup of G. Suppose that for every $x, y \in B$, $x \vee y < \infty \Rightarrow x \vee y \in B$. Then for all $x \in B \cap PP^{-1}$, there exist $\mu, \nu \in B \cap P$ such that $x = \mu\nu^{-1}$ and for all $p, q \in P$ with $pq^{-1} = x$ we have $\mu \leq p$ and $\nu \leq q$.*

The lemma says that if $x \in PP^{-1} \cap B$, then the minimal elements of (2.1) must also be contained in B. In particular, if $\phi(A \cap P) = B \cap P$, then $\phi(x) \in PP^{-1}$.

Proof of Lemma 4.3. Fix $x \in B \cap PP^{-1}$. Say $x = st^{-1}$ with $s, t \in P$. Then $x^{-1}s \in P$ and $x \leq s$. Also $e \leq s$, and so $x \vee e < \infty$. Since $e, x \in B$ we get $x \vee e \in B$. Let $\mu = x \vee e$ and $\nu = x^{-1}(x \vee e)$. Then $\mu\nu^{-1} = x \vee e(x^{-1}(x \vee e))^{-1} = x$.

Fix $p, q \in P$ such that $x = pq^{-1}$. Then $x^{-1}p = (pq^{-1})^{-1}p = q$, and so $x \leq p$. Therefore $\mu = x \vee e \leq p$. Now $\mu^{-1}p \in P$, and then $\nu^{-1}q = \mu^{-1}p \in P$ gives $\nu \leq q$. □

We can now prove Theorem 4.1. Its proof is based on [11, Theorem 2.11], and our presentation is helped by Emily Irwin's treatment of [11, Theorem 2.11] in her University of Otago Honours thesis.

Proof of Theorem 4.1. Fix $x \in P^*P^{*-1}$. We shall prove that there exist $\mu, \nu \in P^*$ with $x = \mu\nu^{-1}$ such that whenever $\gamma\delta^{-1} = x$ we have $\mu \leq \gamma$ and $\nu \leq \delta$ (see (2.1)).

Choose $\alpha, \beta \in P^*$ such that $x = \alpha\beta^{-1}$. Choose a complete set X of left coset representatives of G/A that contains

$$L_A := \{p \in P : q \in pA \cap P \Rightarrow p \leq q\}.$$

By Lemma 4.2 we can write α and β in unique normal form:

$$\alpha = p_0 t p_1 t \dots t p_m t r \text{ where } p_i \in L_A \text{ and } r \in P;$$

$$\beta = q_0 t q_1 t \dots t q_n t s \text{ where } q_i \in L_A \text{ and } s \in P.$$

Now $x = \alpha\beta^{-1}$ is equal to

$$\alpha\beta^{-1} = p_0 t p_1 t \dots t p_m t r s^{-1} t^{-1} q_n^{-1} t^{-1} \dots t^{-1} q_0^{-1}.$$

First we look for initial cancellations in the middle of $\alpha\beta^{-1}$: if $rs^{-1} \in B$, then we can replace $trs^{-1}t^{-1}$ with $\phi^{-1}(rs^{-1})$. By assumption (3), Lemma 4.3 applies and there exist $b_1, b_2 \in P \cap B$ such that $rs^{-1} = b_1 b_2^{-1}$. Then

$$\phi^{-1}(rs^{-1}) = \phi^{-1}(b_1 b_2^{-1}) = \phi^{-1}(b_1)\phi^{-1}(b_2)^{-1}.$$

Since $\phi(A \cap P) = B \cap P$ we have $\phi^{-1}(b_1)\phi^{-1}(b_2)^{-1} \in PP^{-1}$. Then

$$x = \alpha\beta^{-1} = p_0 t p_1 t \dots t p_m \phi^{-1}(b_1)\phi^{-1}(b_2)^{-1} q_n^{-1} t^{-1} \dots t^{-1} q_0^{-1}. \tag{4.3}$$

We can repeat this process until there are no more cancellations available in the middle, and so we assume this is the case for the expression (4.3). This gives the following cases:

(1) there are no t and no more t^{-1},
(2) there are no more t^{-1},
(3) there are no more t,
(4) there are t and t^{-1}, and then the term with t to the left and t^{-1} to its right is not in B.

In each case, we will write down our candidates for μ and ν and prove that they are the required minimums.

(1) Suppose that after initial cancellations, there are no more t and no more t^{-1}. Then $\alpha\beta^{-1} = p_0q_0^{-1}$ is already in normal form. By (2.1) there exist $\sigma, \tau \in P$ such that $p_0q_0^{-1} = \sigma\tau^{-1}$ and for all $c, d \in P$ such that $cd^{-1} = \sigma\tau^{-1}$ we have $\sigma \leq c$ and $\tau \leq d$. So we write $x = \sigma\tau^{-1}$ and choose as our candidates $\mu = \sigma$ and $\nu = \tau$.

Let $\gamma, \delta \in P^*$ such that $x = \gamma\delta^{-1}$. Let θ be the height map. Then $\theta(x) = 0$ and hence $\theta(\gamma) = \theta(\delta)$. We will prove that $\mu \leq \gamma$ and $\nu \leq \delta$ by induction on $\theta(\gamma)$.

For $\theta(\gamma) = 0$ we have $\gamma, \delta \in P$, and then $\mu = \sigma \leq \gamma$ and $\nu = \tau \leq \delta$. Let $k \geq 0$ and suppose that for all $\gamma, \delta \in P^*$ such that $\theta(\gamma) = \theta(\delta) = k$ and $x = \gamma\delta^{-1}$ we have $\mu \leq \gamma$ and $\nu \leq \delta$. Now consider $\gamma, \delta \in P^*$ such that $x = \gamma\delta^{-1}$ and $\theta(\gamma) = \theta(\delta) = k + 1$. We write $\gamma = m_0t \dots m_ktm_{k+1}$ and $\delta = n_0t \dots n_ktn_{k+1}$ in normal form where $m_i, n_i \in L_A$ for $0 \leq i \leq k$ and $m_{k+1}, n_{k+1} \in P$. Next we reduce $x = \gamma\delta^{-1}$ towards normal form. We have

$$x = \gamma\delta^{-1} = m_0t \dots m_ktm_{k+1}n_{k+1}^{-1}t^{-1}n_k \dots tn_0^{-1}.$$

Since x has a unique normal form with no t or t^{-1}, there must be some cancellation. Since the $m_i, n_i \in L_A$ for $0 \leq i \leq k$, the cancellation must occur across $m_{k+1}n_{k+1}^{-1}$. So $m_{k+1}n_{k+1}^{-1} \in B$ and $tm_{k+1}n_{k+1}^{-1}t^{-1} = \phi^{-1}(m_{k+1}n_{k+1}^{-1})$, and

$$\begin{aligned} x = \gamma\delta^{-1} &= m_0t \dots tm_k(tm_{k+1}n_{k+1}^{-1}t^{-1})n_kt^{-1} \dots tn_0^{-1} \\ &= m_0t \dots tm_k\phi^{-1}(m_{k+1}n_{k+1}^{-1})n_kt^{-1} \dots tn_0^{-1} \end{aligned}$$

By assumption (3), Lemma 4.3 applies and there exists $b_m, b_n \in B \cap P$ such that $m_{k+1}n_{k+1}^{-1} = b_mb_n^{-1}$ and $b_m \leq m_{k+1}$ and $b_n \leq n_{k+1}$. Then

$$x = m_0t \dots tm_k\phi^{-1}(b_m)\phi^{-1}(b_n^{-1})n_kt^{-1} \dots tn_0^{-1}.$$

Since $\phi(A \cap P) = B \cap P$ we have $m_{k+1}\phi^{-1}(b_m), n_{k+1}\phi^{-1}(b_n) \in P$. But now we have $\gamma' = m_0t \dots tm_k\phi^{-1}(b_m)$ and $\delta' = n_0t \dots tn_k\phi^{-1}(b_n)$ such that $\gamma'(\delta')^{-1} = x$ and $\theta(\gamma) = \theta(\delta') = k$. By our induction hypothesis we have $\mu \leq \gamma'$ and $\nu \leq \delta'$.

To show that $\mu \leq \gamma$ we compute:

$$\begin{aligned} \gamma &= m_0t \dots tm_ktm_{k+1} \\ &= m_0t \dots tm_ktb_mb_m^{-1}m_{k+1} \\ &= m_0t \dots tm_k\phi^{-1}(b_m)tb_m^{-1}m_{k+1} \quad \text{(replacing } tb_m \text{ with } \phi^{-1}(b_m)t) \\ &= \gamma' tb_m^{-1}m_{k+1}. \end{aligned}$$

Since $b_m^{-1}m_{k+1} \in P$ we have $tb_m^{-1}m_{k+1} \in P^*$. Now $(\gamma')^{-1}\gamma \in P^*$ and $\gamma' \leq \gamma$. Since $\mu \leq \gamma'$ this gives $\mu \leq \gamma$. Similarly, $\delta = \delta' tb_n^{-1}n_{k+1}$ and so $\nu \leq \delta$. By induction, for all $\gamma, \delta \in P^*$ such that $x = \gamma\delta^{-1}$ we have $\mu \leq \gamma$ and $\nu \leq \delta$.

(2) Suppose that after the initial cancellations there are no more t^{-1} left. Then we have x in normal form:

$$x = \alpha\beta^{-1} = p_0tp_1t \dots tp_{m-n-1}trs^{-1}$$

where $m \geq n$. By (2.1) there exist $\sigma, \tau \in P$ such that $rs^{-1} = \sigma\tau^{-1}$ and $\sigma \leq r$ and $\tau \leq s$. So

$$x = p_0tp_1t \dots tp_{m-n-1}t\sigma\tau^{-1}. \tag{4.4}$$

Our candidates are

$$\mu = p_0 t p_1 t \dots t p_{m-n-1} t \sigma \quad \text{and} \quad \nu = \tau.$$

Fix $\gamma, \delta \in P^*$ such that $x = \gamma\delta^{-1}$. Say $\gamma = m_0 t m_1 \dots t m_i$ and $\delta = n_0 t n_1 \dots n_j$ in normal form. Then

$$\gamma\delta^{-1} = m_0 t m_1 \dots t m_i n_j^{-1} t^{-1} \dots n_1 t^{-1} n_0.$$

Since $\theta(\gamma\delta^{-1}) = \theta(\mu\nu^{-1}) = m - n$ we get

$$i = \theta(\gamma) = \theta(\delta) + m - n$$

and hence $i \geq m - n$. It follows from the uniqueness of normal form that there exists $\gamma' \in P^*$ such that

$$\gamma\delta^{-1} = p_0 t p_1 t \dots t p_{m-n-1} t \gamma' \delta^{-1}.$$

Thus $p_0 t p_1 t \dots t p_{m-n-1} t \leq \gamma$. From (4.4) we have $(p_0 t p_1 t \dots t p_{m-n-1} t)^{-1} x = \sigma\tau^{-1}$. Since $(p_0 t p_1 t \dots t p_{m-n-1} t)^{-1}\gamma, \delta \in P^*$ and $(p_0 t p_1 t \dots t p_{m-n-1} t)^{-1}\gamma\delta = \sigma\tau^{-1}$, we can apply (1) above with $\mu' = \sigma$ and $\nu' = \tau$ to see that $\sigma \leq (p_0 t p_1 t \dots t p_{m-n-1} t)^{-1}\gamma$ and $\tau \leq \delta$. Hence $\mu = p_0 t p_1 t \dots t p_{m-n-1} t\sigma \leq \gamma$ and $\nu \leq \delta$ as required.

(3) Suppose that after the cancellations, there are no more t left. Then

$$x = r s^{-1} t^{-1} q_{n-m-1}^{-1} t^{-1} \dots t^{-1} q_0^{-1}$$

for some $r, s \in P$. Consider

$$x^{-1} = q_0 t q_1 t \dots t q_{n-m-1} t s r^{-1}.$$

By (2.1) there exist $\sigma, \tau \in P$ such that $rs^{-1} = \sigma\tau^{-1}$ and $\sigma \leq r$ and $\tau \leq s$. Let

$$\mu = \sigma \quad \text{and} \quad \nu = q_0 t q_1 t \dots t q_{n-m-1} t \tau.$$

By (2) they have the property that $x^{-1} = \nu\mu^{-1}$ and for all $\gamma, \delta \in P^*$ such that $x^{-1} = \delta\gamma^{-1}$ we have $\mu \leq \gamma$ and $\nu \leq \delta$. Taking inverses, $x = \mu\nu^{-1}$ and for all $\gamma, \delta \in P^*$ such that $x = \gamma\delta^{-1}$ we have $\mu \leq \gamma$ and $\nu \leq \delta$.

(4) Suppose that after the initial cancellations there are both t and t^{-1} left. Then the term with t to the left and t^{-1} to its right is not in B. There exist $k \leq m$ and $l \leq n$ such that

$$x = p_0 t p_1 t \dots t p_k t r s^{-1} t^{-1} q_l^{-1} t^{-1} \dots t^{-1} q_0^{-1}.$$

By (2.1) there exist $\sigma, \tau \in P$ such that $rs^{-1} = \sigma\tau^{-1}$ and $\sigma \leq r$ and $\tau \leq s$. Our candidate for μ, ν are

$$\mu = p_0 t p_1 t \dots t p_k t \sigma \quad \text{and} \quad \nu = q_0 t q_1 t \dots t q_l t \tau.$$

Fix $\gamma, \delta \in P^*$ such that $x = \gamma\delta^{-1}$. By the argument used in (2), there exists $\gamma' \in P^*$ such that $\gamma\delta^{-1} = p_0 t p_1 t \dots t p_j t \gamma' \delta^{-1}$. Hence $\gamma' = (p_0 t p_1 t \dots t p_j t)^{-1}\gamma \in P^*$ and $p_0 t p_1 t \dots t p_j t \leq \gamma$.

Consider $\gamma'\delta^{-1} = \sigma\tau^{-1} t^{-1} q_k^{-1} t^{-1} \dots t^{-1} q_0^{-1}$. Here $\gamma', \delta \in P^*$ and there are no t in $\gamma'\delta^{-1}$ after cancellation. Applying (3) with $\mu' = \sigma$ and $\nu' = \nu$ to get $\mu' = \sigma \leq \gamma'$ and $\nu' = \nu \leq \delta$. Then $\mu = p_0 t p_1 t \dots t p_j t\sigma \leq p_0 t p_1 t \dots t p_j t \gamma' = \gamma$. □

Theorem 4.1 gives new examples of quasi-lattice ordered HNN extensions.

Example 4.4. We can show that the Baumslag-Solitar group $(\mathrm{BS}(c,d), \mathrm{BS}(c,d)^+)$ with $c, d > 0$ is quasi-lattice ordered using Theorem 4.1. Since $(\mathbb{Z}, \mathbb{N})$ is totally ordered it is quasi-lattice ordered. Let $A = \{dm : m \in \mathbb{Z}\}$ and $B = \{cm : m \in \mathbb{Z}\}$. Every element $n \in \mathbb{N}$ has a unique decomposition $n = r + md$ where $m \in \mathbb{N}$ and $r \in \{0, 1, \dots, d-1\}$. The remainder r is a choice of coset representative $n + A = r + A$. For all $n' \in n + A \cap \mathbb{N}$ we have $n' = r + md + kd$ where $k \in \mathbb{Z}$ and $k \leq m$. Thus $r \leq n'$. Hence every coset of $\mathbb{Z}/A$ has nontrivial intersection with $\mathbb{N}$, and has a minimal coset representative in $\mathbb{N}$. Since B is totally ordered it is closed under taking least upper bounds. Define $\phi : A \to B$ by $\phi(dm) = cm$. Then $\phi(A \cap \mathbb{N}) = B \cap \mathbb{N}$. So Theorem 4.1 applies and gives that $(\mathbb{Z}^*, \mathbb{N}^*)$ is quasi-lattice ordered.

Example 4.5. We can generalise the previous example to $(\mathbb{Z}^2, \mathbb{N}^2)$, which is quasi-lattice ordered by [9, Example 2.3(2)]. Fix $a, b, c, d \in \mathbb{N}$. Then $A = \{(am, bn) : m, n \in \mathbb{Z}\}$ and $B = \{(cm, dn) : m, n \in \mathbb{Z}\}$ are subgroups of $\mathbb{N}^2$. Let $\phi : A \to B$ be defined by $\phi((am, bn)) = (cm, dn)$. This ϕ satisfies $\phi(A \cap \mathbb{N}^2) = B \cap \mathbb{N}^2$. For all $(m, n) \in \mathbb{N}^2$, the division algorithm on $\mathbb{N}$ gives a unique decomposition

$$(m, n) = (r_1, r_2) + (ja_1, ka_2) \text{ for } j, k \in \mathbb{N} \text{ and } r_1 \in \{0, \dots a_1 - 1\},\ r_2 \in \{0, \dots, a_2 - 1\}.$$

Thus (r_1, r_2) is a minimal left coset representative of $(m, n) + A$. For all $(m, n), (p, q) \in B$, we have

$$(m, n) \vee (p, q) = (\max\{m, p, 0\}, \max\{n, q, 0\})$$

and hence $(m, n) \vee (p, q) \in B$. So B is closed under $\vee$. By Theorem 4.1, $(\mathbb{Z}^{2*}, \mathbb{N}^{2*})$ is a quasi-lattice ordered group with presentation

$$\mathbb{Z}^{2*} = \langle \mathbb{Z}^2 \cup \{t\} \mid (am, bn)t = t(cm, dn) \rangle.$$

It is straightforward to extend this construction to $(\mathbb{Z}^n, \mathbb{N}^n)$.

Example 4.6. Consider the free group $\mathbb{F}_2$ on 2 generators $\{a, b\}$ and let $\mathbb{F}_2^+$ be the subsemigroup generated by e, a and b. The pair $(\mathbb{F}_2, \mathbb{F}_2^+)$ is quasi-lattice ordered by [9, Example 2.3(4)]. Let $A = \{a^n : n \in \mathbb{Z}\}$, $B = \{b^n : n \in \mathbb{Z}\}$ and $\phi : A \to B$ defined by $\phi(a^n) = b^n$. Every $x \in F_2^+$ can be written as a product of $y \in \mathbb{F}_2^+$ which does not end in a followed by a^n for some $n \geq 0$. Then $y \in xA$. Every $z \in yA \cap \mathbb{F}_2^+$ begins with the word y which is in $\mathbb{F}_2^+$. It follows that $y \leq z$. Thus A has minimal left coset representatives in $\mathbb{F}_2^+$. Since B is totally ordered, it is trivially closed under $\vee$. It follows from Theorem 4.1 that $(\mathbb{F}_2^*, \mathbb{F}_2^{+*})$ is a quasi-lattice ordered group with presentation

$$\mathbb{F}_2^* = \langle \{a, b, t\} \mid at = tb \rangle.$$

Example 4.7. Building on $(\mathbb{F}_2, \mathbb{F}_2^+)$ again, fix $s, u \in \mathbb{N}$, and let $A = \{a^{ms} : m \in \mathbb{Z}\}$, $B = \{b^{mu} : m \in \mathbb{Z}\}$ and $\phi : A \to B$ be $\phi(a^{ms}) = b^{mu}$. Then B is totally ordered and hence is closed under $\vee$. To see that A has minimal coset representatives, we observe that every $x \in \mathbb{F}_2^+$ is a product of a $y \in \mathbb{F}_2^+$ that does not end in a followed by a^n for some $n \in \mathbb{N}$. We write $n = r + js$ for some $j \in \mathbb{N}$ and $r \in \{0, 1, \dots, s-1\}$. We choose ya^r as our coset representative. Then for all $z \in ya^r A \cap \mathbb{F}_2^+$ we have $ya^r \leq z$. It follows from Theorem 4.1 that $(\mathbb{F}_2^*, \mathbb{F}_2^{+*})$ is a quasi-lattice ordered group with presentation

$$\mathbb{F}_2^* = \langle \{a, b, t\} \mid a^s t = tb^u \rangle.$$

Taking $u = s = 1$ gives Example 4.6.

Replacing B by $B' = \{a^{mu} : m \in \mathbb{Z}\}$ gives a quasi-lattice ordered group $(\mathbb{F}_2^*, \mathbb{F}_2^{+*})$ with presentation

$$\mathbb{F}_2^* = \langle \{a, b, t\} \mid a^s t = t a^u \rangle.$$

In the next two examples we show that it is easy to find subgroups which do not have minimal left coset representatives.

Example 4.8. Consider the group

$$G = \mathbb{Z}(\sqrt{2}) = \{m + n\sqrt{2} : m, n \in \mathbb{Z}\}$$

with subsemigroup $\mathbb{Z}(\sqrt{2})^+ = \mathbb{Z}(\sqrt{2}) \cap [0, \infty)$. Let $A = \mathbb{Z}(2\sqrt{2})$. We claim there are no minimal coset representatives for A. Suppose, aiming for a contradiction, that there exists some coset representative $p \in \mathbb{Z}(\sqrt{2})^+$ such that

$$q \in [p + \mathbb{Z}(2\sqrt{2})] \cap \mathbb{Z}(\sqrt{2})^+ \Rightarrow p \leq q.$$

Recall that $\mathbb{Z}(2\sqrt{2})$ is dense in $\mathbb{R}$.[2] Thus there exists some $a \in \mathbb{Z}(2\sqrt{2}) \cap (0, p)$ hence $a \leq p$. Thus $p - a \in [p + \mathbb{Z}(2\sqrt{2})] \cap \mathbb{Z}(\sqrt{2})^+$. But $p - a \leq p$, giving a contradiction.

Example 4.9. Consider $(\mathbb{Z}^2, \mathbb{N}^2)$, and let A be the subgroup generated by $\{(1, 2), (2, 1)\}$. Consider the coset

$$(1, 0) + A = (0, 1) + A.$$

Since $(1, 0)$ and $(0, 1)$ have no nonzero lower bound, there can be no choice of minimal coset representative.

5. Amenability of (G^*, P^*)

In this section we prove the following theorem.

Theorem 5.1. *Let (G, P) be a quasi-lattice ordered group with subgroups A and B. Suppose that $\phi : A \to B$ is an isomorphism which satisfies the hypotheses of Theorem 4.1. Let (G^*, P^*) be the corresponding HNN extension. If (G, P) is amenable, then (G^*, P^*) is amenable.*

To prove the theorem we will show that the height map $\theta : G^* \to \mathbb{Z}$ is a controlled map, that $(\theta^{-1}(e), \theta^{-1}(e) \cap P^*)$ is amenable, and then apply Theorem 3.2. To prove that $(\theta^{-1}(e), \theta^{-1}(e) \cap P^*)$ is amenable, we start by investigating order-preserving isomorphisms between the semigroups of quasi-lattice ordered groups.

Lemma 5.2. *Let (G, P) and (K, Q) be quasi-lattice ordered groups. Suppose that there is a semigroup isomorphism $\phi : P \to Q$. Then ϕ is order-preserving. In particular, for $x, y \in P$, $x \vee y < \infty$ if and only if $\phi(x) \vee \phi(y) < \infty$. If $x \vee y < \infty$ then $\phi(x \vee y) = \phi(x) \vee \phi(y)$.*

Proof. To see that ϕ is order-preserving, let $x, y \in P$ and $x \leq y$. Then $y \in xP$, and $\phi(y) \in \phi(x)\phi(P) = \phi(x)Q$. Thus $\phi(x) \leq \phi(y)$, and ϕ is order-preserving.

To show that ϕ preserves the least upper bound structure, first suppose that $x, y \in P$ such that $x \vee y < \infty$. Since ϕ is order-preserving it follows that $\phi(x), \phi(y) \leq \phi(x \vee y)$.

[2]To see denseness observe that $0 < (-2 + 2\sqrt{2}) < 1$ and $(-2 + 2\sqrt{2})^n \in \mathbb{Z}(\sqrt{2})$ for all $n \in \mathbb{N}$. Thus for every open interval (u, v) there exists n such that $(-2 + 2\sqrt{2})^n < v - u$. Hence there exists $k \in \mathbb{N}$ such that $k(-2 + 2\sqrt{2})^n \in (u, v)$.

Thus $\phi(x), \phi(y)$ have a common upper bound in Q. Hence $\phi(x) \vee \phi(y)$ exists and $\phi(x) \vee \phi(y) \leq \phi(x \vee y)$. Second, suppose that $\phi(x) \vee \phi(y) < \infty$ for some $x, y \in P$. Since $\phi^{-1} : Q \to P$ is a semigroup isomorphism it is order-preserving. Thus $\phi^{-1}(\phi(x) \vee \phi(y))$ is an upper bound for $x = \phi^{-1}(\phi(x))$ and $y = \phi^{-1}(\phi(y))$. Hence $x \vee y$ exists and $x \vee y \leq \phi^{-1}(\phi(x) \vee \phi(y))$. Thus

$$\phi(x \vee y) \leq \phi \circ \phi^{-1}(\phi(x) \vee \phi(y)) = \phi(x) \vee \phi(y).$$

Hence $x \vee y < \infty$ if and only if $\phi(x) \vee \phi(y) < \infty$, and $\phi(x \vee y) = \phi(x) \vee \phi(y)$. □

Proposition 5.3. *Let (G, P) and (K, Q) be quasi-lattice ordered groups. Let $\{v_p : p \in P\}$ and $\{w_q : q \in Q\}$ be the generating elements of $C^*(G, P)$ and $C^*(K, Q)$, respectively. Suppose that there is a semigroup isomorphism $\phi : P \to Q$.*

1. *There exists an isomorphism $\pi_\phi : C^*(G, P) \to C^*(K, Q)$ such that $\pi_\phi(v_p) = w_{\phi(p)}$.*
2. *(G, P) is amenable if and only if (K, Q) is amenable.*

Proof. (1) We will show that $T : P \to C^*(K, Q)$ defined by $T_p = w_{\phi(p)}$ is a covariant representation of P, and then take $\pi_\phi := \pi_T$. Fix $p, q \in P$. Since ϕ is a semigroup isomorphism we have

$$T_p T_q = w_{\phi(p)} w_{\phi(q)} = w_{\phi(pq)} = T_{pq}$$

and $T_e = w_{\phi(e_G)} = w_{e_K} = 1$. Hence T is an isometric representation. We have

$$\begin{aligned} T_p T_p^* T_q T_q^* &= w_{\phi(p)} w_{\phi(p)}^* w_{\phi(q)} w_{\phi(q)}^* \\ &= \begin{cases} w_{\phi(p)\vee\phi(q)} w_{\phi(p)\vee\phi(q)}^* & \text{if } \phi(p) \vee \phi(q) < \infty \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

which, using Lemma 5.2, is equivalent to

$$\begin{aligned} &= \begin{cases} w_{\phi(p\vee q)} w_{\phi(p\vee q)}^* & \text{if } p \vee q < \infty \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} T_{p\vee q} T_{p\vee q}^* & \text{if } p \vee q < \infty \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Hence T is covariant. By the universal property of $C^*(G, P)$ there exists a homomorphism $\pi_\phi : C^*(G, P) \to C^*(K, Q)$ such that $\pi_\phi(v_p) = T_p = w_{\phi(p)}$.

Since $\phi^{-1} : Q \to P$ is an isomorphism the argument above gives a homomorphism $\pi_{\phi^{-1}} : C^*(K, Q) \to C^*(G, P)$ such that $\pi_{\phi^{-1}}(w_q) = v_{\phi^{-1}(q)}$. In particular,

$$\pi_{\phi^{-1}}(\pi_\phi(v_p)) = \pi_{\phi^{-1}}(w_{\phi(p)}) = v_{\phi^{-1}(\phi(p))} = v_p$$

and $\pi_\phi(\pi_{\phi^{-1}}(w_q)) = w_q$ It follows that π_ϕ is an isomorphism from $C^*(G, P)$ to $C^*(K, Q)$.

(2) By symmetry it suffices to show that if (K, Q) is amenable then (G, P) is amenable. Let E_Q and E_P be the conditional expectations on $C^*(K, Q)$ and $C^*(G, P)$, respectively. To see $E_P = \pi_\phi^{-1} \circ E_Q \circ \pi_\phi$ we compute on spanning elements:

$$\pi_\phi^{-1} \circ E_Q \circ \pi_\phi(v_p v_q^*) = \pi_\phi^{-1} \circ E_Q(w_{\phi(p)} w_{\phi(q)}^*)$$

$$= \begin{cases} \pi_\phi^{-1}(w_{\phi(p)}w^*_{\phi(q)}) & \text{if } \phi(p) = \phi(q) \\ 0 & \text{otherwise} \end{cases}$$
$$= \begin{cases} \pi_\phi^{-1}(w_{\phi(p)}w^*_{\phi(q)}) & \text{if } p = q \\ 0 & \text{otherwise} \end{cases}$$
$$= \begin{cases} v_p v_q^* & \text{if } p = q \\ 0 & \text{otherwise} \end{cases}$$
$$= E_P(v_p v_q^*).$$

Suppose that (K, Q) is amenable. Then E_Q is faithful. Suppose that $a \geq 0$ and $E_P(a) = 0$. Then $\pi_\phi(a) \geq 0$, and

$$0 = E_P(a) = \pi_\phi^{-1} \circ E_Q \circ \pi_\phi(a) \Rightarrow 0 = E_Q \circ \pi_\phi(a) \Rightarrow 0 = \pi_\phi(a)$$

because E_Q is faithful. Since π_ϕ is faithful, $a = 0$. Now E_P is faithful, and hence (G, P) is amenable. □

Next we need some lemmas which will be used to show that the height map θ is a controlled map. In particular we need to identify the minimal elements of Definition 3.1. If $x \in P^*$ has normal form

$$x = p_0 t p_1 t \dots p_{n-1} t p_n$$

we call $p_0 t p_1 t \dots p_{n-1} t$ the *stem* of x and write

$$\text{stem}(x) = p_0 t p_1 t \dots p_{n-1} t.$$

The set of stems is our candidate for the minimal elements.

Lemma 5.4. *Let (G, P) be a quasi-lattice ordered group with subgroups A and B. Suppose that $\phi : A \to B$ is an isomorphism which satisfies the hypotheses of Theorem 4.1. Let $p, q \in P$. Then p and q have a common upper bound in P^* if and only if p and q have a common upper bound in P.*

Proof. First suppose that p and q have a common upper bound $r \in P$. Then $r \in P^*$ and so r is a common upper bound for p and q in P^*.

Second, suppose that p and q have a common upper bound $x \in P^*$. If $\theta(x) = 0$, then $x \in P$ and we are done. Suppose, aiming for a contradiction, that $\theta(x) = k$ for some $k \geq 1$, and that p, q have no common upper bound y with $\theta(y) < k$.

Observe that $p^{-1}x, q^{-1}x \in P^*$, and that $\theta(p^{-1}x) = \theta(x) = \theta(q^{-1}x) = k$. We write $p^{-1}x$ and $q^{-1}x$ in their normal forms:

$$p^{-1}x = p_0 t p_1 t \dots p_{k-1} t p_k \quad \text{and} \quad q^{-1}x = q_0 t q_1 t \dots q_{k-1} t q_k,$$

where $p_i, q_i \in L_A$ for $i < k$ and $p_k, q_k \in P$. Consider

$$p^{-1}q = p^{-1}x(q^{-1}x)^{-1} = p_0 t p_1 t \dots p_{k-1} t p_k q_k^{-1} t^{-1} q_{k-1}^{-1} \dots t^{-1} q_0^{-1}.$$

Since $p^{-1}x(q^{-1}x)^{-1} = p^{-1}q$ and $p^{-1}q$ is in normal form we must have some cancellation. Since the first k terms are already in normal form, $p_k q_k^{-1} \in B$. By Lemma 4.3, there exist $b_1, b_2 \in B \cap P$ such that $b_1 \leq p_k$, $b_2 \leq q_k$ and $b_1 b_2^{-1} = p_k q_k^{-1}$. Then

$$p^{-1}q = p^{-1}x(q^{-1}x)^{-1} = p_0 t \dots p_{k-1} t (p_k q_k^{-1}) t^{-1} q_{k-1}^{-1} \dots t^{-1} q_0^{-1}$$

$$\begin{aligned}
&= p_0 t \dots p_{k-1} t (b_1 b_2^{-1}) t^{-1} q_{k-1}^{-1} \dots t^{-1} q_0^{-1} \\
&= p_0 t \dots p_{k-1} \phi^{-1}(b_1) t t^{-1} \phi^{-1}(b_2)^{-1} q_{k-1}^{-1} \dots t^{-1} q_0^{-1} \\
&= p_0 t \dots p_{k-1} \phi^{-1}(b_1) \phi^{-1}(b_2)^{-1} q_{k-1}^{-1} \dots t^{-1} q_0^{-1}.
\end{aligned}$$

Rearranging gives

$$p(p_0 t \dots t p_{k-1} \phi^{-1}(b_1)) = q(q_0 t q_1 t \dots t q_{k-1} \phi^{-1}(b_2)) \in P^*.$$

Therefore $y = p(p_0 t p_1 t \dots p_{k-1} \phi^{-1}(b_1))$ is a common upper bound for p and q in P^* and $\theta(y) = k-1$, giving us the contradiction we sought. Therefore p and q have a common upper bound y with $\theta(y) = 0$, and hence they have a common upper bound in P. □

The statement of Lemma 5.5 is adapted from [2, Lemma 3.4].

Lemma 5.5. *Let $x, y \in P^*$ such that $x \vee y < \infty$. Write*

$$x = \mathrm{stem}(x)p \text{ and } y = \mathrm{stem}(y)q \text{ where } p, q \in P.$$

(1) *If $\theta(x) = \theta(y)$, then $\mathrm{stem}(x) = \mathrm{stem}(y)$ and $p \vee q < \infty$. In particular, $x \vee y = \mathrm{stem}(x)(p \vee q)$ and $\theta(x \vee y) = \theta(x) = \theta(y)$.*
(2) *If $\theta(x) < \theta(y)$, then there exists $r \in P$ such that $x \vee y = yr$ and $\theta(x \vee y) = \theta(y)$.*

In particular, $\theta(x \vee y) = \max\{\theta(x), \theta(y)\}$.

Proof. (1) Suppose that $\theta(x) = \theta(y)$. We know that $x \le x \vee y$ and $y \le x \vee y$. Thus, by the uniqueness of normal forms, $\mathrm{stem}(x) = \mathrm{stem}(y)$. Now by left invariance of the partial order we see that

$$p = \mathrm{stem}(x)^{-1} x \le \mathrm{stem}(x)^{-1}(x \vee y) \quad \text{and} \quad q = \mathrm{stem}(x)^{-1} y \le \mathrm{stem}(x)^{-1}(x \vee y).$$

Therefore p and q have a common left upper bound in P^* and hence, by Lemma 5.4, they have a common left upper bound in P and $p \vee q$ exists in P. By left invariance $x \vee y = \mathrm{stem}(x)p \vee q$. Further, $\theta(x \vee y) = \theta(x) = \theta(y)$.

(2) Suppose that $\theta(x) < \theta(y)$. Since $x \le x \vee y$ we have $x^{-1}(x \vee y) \in P^*$. We can write $x^{-1}(x \vee y) = \tau\gamma u$ for some $u \in P$, $\tau \in \Sigma_{\theta(y)-\theta(x)}$ and $\gamma \in \Sigma_{\theta(x\vee y)-\theta(y)}$. Then $x \vee y = x\tau\gamma u$.

Now we have $x\tau \le x \vee y$ and $\theta(x\tau) = \theta(x) + (\theta(y) - \theta(x)) = \theta(y)$. Write $x\tau = \mathrm{stem}(x\tau)w$ for some $w \in P$. Therefore $x\tau \vee y < \infty$ and $\theta(x\tau) = \theta(y)$ so we can apply (1) to see that $\mathrm{stem}(x\tau) = \mathrm{stem}(y)$ and $x\tau \vee y = \mathrm{stem}(y)(q \vee w)$.

Now $x \vee y \le \mathrm{stem}(y)(q \vee w)$. Therefore there exists some $r \in P$ such that $x \vee y = \mathrm{stem}(y)qr = yr$. Then $\theta(x \vee y) = \theta(y)$.

By (1) and (2) we see that

$$\theta(x \vee y) = \begin{cases} \theta(x) & \text{if } \theta(x) = \theta(y) \\ \theta(y) & \text{if } \theta(x) < \theta(y). \end{cases}$$

Thus $\theta(x \vee y) = \max\{\theta(x), \theta(y)\}$. □

Proof of Theorem 5.1. We will use Theorem 3.2; to do so we need to show that the height map $\theta : (G^*, P^*) \to (\mathbb{Z}, \mathbb{N})$ is a controlled map in the sense of Definition 3.1, and that $(\theta^{-1}(e), \theta^{-1}(e) \cap P^*)$ is amenable.

To see that θ is order-preserving, let $x, y \in P^*$ such that $x \leq y$. Then $x^{-1}y \in P^*$ and $\theta(x^{-1}y) \geq 0$. So $0 \leq \theta(x^{-1}y) = -\theta(x) + \theta(y)$ and hence $\theta(x) \leq \theta(y)$. By Lemma 5.5, if $x \vee y < \infty$, then $\theta(x \vee y) = \max\{\theta(x), \theta(y)\} = \theta(x) \vee \theta(y)$.

For every $k \in \mathbb{N}$, Σ_k is complete: if $x \in \theta^{-1}(k) \cap P^*$, then $\mathrm{stem}(x) \in \Sigma_k$ and $x = \mathrm{stem}(x)p$ for some $p \in P$. Hence $\mathrm{stem}(x) \leq x$. By the uniqueness of normal forms, if $\sigma, \tau \in \Sigma_k$ and $\sigma \vee \tau < \infty$ then $\sigma = \tau$. Therefore θ is a controlled map into the amenable group $\mathbb{Z}$.

Suppose that (G, P) is amenable. Then $\theta^{-1}(0) \cap P^*$ is the set of elements of P^* with height 0, and hence they all have normal form p_0 for some $p_0 \in P$. Thus $\theta^{-1}(0) \cap P^*$ is isomorphic to P. Since (G, P) is amenable, so is $(\theta^{-1}(0), \theta^{-1}(0) \cap P^*)$ by Lemma 5.3. Since $(\mathbb{Z}, \mathbb{N})$ is amenable, it now follows from Theorem 3.2 that (G^*, P^*) is an amenable quasi-lattice ordered group. □

Example 5.6. Since $(\mathbb{Z}^2, \mathbb{N}^2)$ and $(\mathbb{F}_2, \mathbb{F}_2^+)$ are amenable quasi-lattice ordered groups [9, §5.1], Theorem 5.1 implies that the HNN extensions $(\mathbb{Z}^{2*}, \mathbb{N}^{2*})$ and $(\mathbb{F}_2^*, \mathbb{F}_2^{+*})$ in Examples 4.5-4.7 are amenable quasi-lattice ordered groups.

References

1. N. Brownlowe, N.S. Larsen, and N. Stammeier, *On C^*-algebras associated to right LCM semigroups*, Trans. Amer. Math. Soc. **369** (2017), 31–68.
2. L.O. Clark, A. an Huef, and I. Raeburn, *Phase transitions of the Toeplitz algebras of Baumslag-Solitar semigroups*, Indiana Univ. Math. J. **65** (2016), 2137–2173.
3. J. Crisp and M. Laca, *On the Toeplitz algebras of right-angled and finite-type Artin groups*, J. Austral. Math. Soc. **72** (2002), 223–245.
4. J. Crisp and M. Laca, *Boundary quotients and ideals of Toeplitz C^*-algebras of Artin groups*, J. Funct. Anal. **242** (2007), 127–156.
5. M. Laca and I. Raeburn, *Semigroup crossed products and the Toeplitz algebras of nonabelian groups*, J. Funct. Anal. **139** (1996), 415–440.
6. M. Laca and I. Raeburn, *Phase transition on the Toeplitz algebra of the affine semigroup over the natural numbers*, Adv. Math. **225** (2010), 643–688.
7. X. Li, *Semigroup C^*-algebras and amenability of semigroups*, J. Funct. Anal. **262** (2012), 4302–4340.
8. R.C. Lyndon and P.E. Schupp, Combinatorial Group Theory, Springer-Verlag, 1977.
9. A. Nica, *C^*-algebras generated by isometries and Wiener-Hopf operators*, J. Operator Theory **27** (1992), 17–52.
10. I. Raeburn and D.P. Williams, Morita Equivalence and Continuous-Trace C^*-Algebras, Mathematical Surveys and Monographs, vol. 60, Amer. Math. Soc., 1998.
11. J. Spielberg, *C^*-algebras for categories of paths associated to the Baumslag-Solitar groups*, J. Lond. Math. Soc. **86** (2012), 728–754.
12. J. Spielberg, *Groupoids and C^*-algebras for categories of paths*, Trans. Amer. Math. Soc. **366** (2014), 5771–5819.
13. C. Starling, *Boundary quotients of C^*-algebras of right LCM semigroups*, J. Funct. Anal. **268** (2015), 3326–3356.
14. J. Tomiyama, *On the projection of norm 1 on W^*-algebras*, Proc. Japan Acad. **33** (1957), 608–612.

Department of Mathematics and Statistics, University of Otago, PO Box 56, Dunedin 9054, New Zealand.

E-mail address: {astrid, iraeburn, itolich}@maths.otago.ac.nz